LAURUS MAJOR Roſtro inæquali.

The Cut Water.

THE Bill, which is the characteriſtick note of this Bird, is a wonderful work of Nature. The baſis of the upper Mandible is thick and compreſſed ſideways gradually to the end, and terminates in a point, and is three inches long. The under Mandible is more compreſſed than the upper, and very thin, both edges being as ſharp as a knife, and is almoſt an inch longer than the upper Mandible, which has a narrow Grove or Channel, into which the upper Edge of the lower Mandible ſhuts. Half the Bill, next the Head, is red, the reſt is black; the forepart of the Head, Neck, Breaſt and Belly white; the hindpart of the Head, Back and Wings black, with a ſmall Mixture of white. The upper Feather of the Tail is black, the reſt white. The Legs are ſhort and ſmall, of a red Colour. The Feet are webbed like thoſe of a Gull, with a ſmall back Toe. Theſe Birds frequent near the Sea Coaſts of *Carolina*. They fly cloſe to the Surface of the Water, from which they ſeem to receive ſomewhat of Food. They alſo frequent Oyſter-banks, on which, I believe, they feed. The ſtructure of their Bills ſeem adapted to that purpoſe. The Cocks and Hens are alike in colour.

Le Coupeur d'eau.

SON bec, qui eſt la marque characteriſtique de cet oiſeau, eſt un ſurprenant ouvrage de la nature. La mandibule ſuperieure eſt épaiſſe à ſa baſe, applatie par les côtés graduellement juſques à ſon extremité; elle a trois pouces de long, & ſe termine en pointe. La mandibule inferieure eſt plus applatie que la ſuperieure, & fort mince; ſes deux bords ſont auſſi coupants qu'un couteau; elle eſt preſque d'un pouce plus longue que l'autre, qui a un canal fort étroit, dans lequel le bord ſuperieur de la mandibule inferieure entre en ſe fermant. La moitié du bec vers la tête eſt rouge, le reſte eſt noir. Le devant de la tête, le col, & le ventre ſont blancs; le derriere de la tête, le dos & les ailes ſont noirs, avec quelque mélange de blanc. La plume de deſſus de ſa queüe eſt noire, les autres ſont blanches. Ses jambes ſont courtes, petites & rouges: ſes pieds ſont garnis de membranes comme ceux de la Moüette, avec un petit ergot derriere. Ces oiſeaux frequentent proche les côtes de la Coroline. Ils volent tout près de la ſurface de l'eau, d'où il ſemble qu'ils tirent quelque nourriture. Ils frêquentent auſſi les bancs où il y a des Huitres, & je croi qu'ils en mangent. La forme de leur bec paroît propre à cela. Le mâle & la femelle ont les mêmes couleurs.